

Alien and invasive freshwater fish species in Iran: Diversity, environmental impacts and management

Hamid Reza ESMAEILI^{1*}, Azad TEIMORI², Feridon OWFI³, Keivan ABBASI⁴, Brian W. COAD⁵

¹*Ichthyology Research Laboratory, Department of Biology, College of Sciences, Shiraz University, Shiraz, Iran.*

²*Department of Biology, Faculty of Sciences, Shahid-Bahonar University of Kerman, Iran.*

³*Iranian Fishery Research Organization (IFRO), Marine Ecology Department, Tehran, Iran.*

⁴*Aquatic Ecology Department of the Inland Water Aquaculture Center, P.O. Box 66, Bandar-e Anzali, Iran.*

⁵*Canadian Museum of Nature, Ottawa, Ontario, K1P 6P4 Canada.*

Email: hresmaeili22@gmail.com

Abstract: More than 32 alien (exotic) fish species belonging to 10 orders and 12 families (Cyprinidae, Gobiidae, Salmonidae, Anguillidae, Mugilidae, Centrarchidae, Heteropneustidae, Gasterosteidae, Cichlidae, Poeciliidae, Adrianichthyidae and Pleuronectidae) were reported from Iran of which 25 species are confirmed by specimens. Some of the alien fishes (e.g., *Cyprinus carpio*, *Carassius auratus*, *Pseudorasbora parva*, *Xiphophorus hellerii* and *Gambusia holbrooki*) have already established breeding populations, acting as invasive species. Some others are regularly stocked by the Iranian Fisheries Company (e.g., *Hypophthalmichthys molitrix*, *Hypophthalmichthys nobilis*, *Ctenopharyngodon idella*) and few others are occasionally recorded from natural waters. *Amatitlania nigrofasciata* and *Tilapia zillii* are two alien cichlids recently reported from Iran. Aquaculture, sport fishing, control of malaria, ornamental purposes, research activities, demonstration in national fairs and accidental introduction are the main reasons for these introductions. *Pseudorasbora parva* was introduced accidentally along with other exotic carps and now is widely distributed in inland waters. The intentional introduction of alien species is an extraordinarily complex issue and requires taking into account a broad number of variables. Understanding the risks that alien species, especially aquatic invasive species (AIS) pose, comparative studies on the ecology and life history strategies of the introduced and invasive species and closely related indigenous species with similar ecological life traits, monitoring the rate of invasion, and the causal relation between population dynamics and species invasion and public awareness, could be effective management strategies to minimize the bio-invasion impacts.

Keywords: Exotic, Threats, Diversity, Management, Iran.

Introduction

Invasive species (IS) are alien (non-native) organisms that have been introduced into an area outside of their natural range, establishing self-sustaining populations and spreading beyond their initial point of introduction, with deleterious impacts on the environment, economy and human health (Kolar & Lodge 2001; Lymbery et al. 2014). Biological invasions are now considered a major environmental issue of public concern (Gozlan et al.

2010). Human population growth, increasing transport capacity and economic globalisation have accelerated the rate of introductions of alien species throughout the world (Vitousek et al. 1997; Sakai et al. 2001; Lymbery et al. 2014). Aquatic invasive species (AIS) and bio-invasions are global environmental issues in marine, brackish and freshwater ecosystems of the world. AIS have been moved across the globe and introduced in totally new environments as an effort to enhance, restore or re-

Table 1. Terminology for alien species (after Lymbery et al. (2014) and Gozlan et al. (2010)).

Term	Definition
Native species	A species occurring within the range it occupies (or could occupy) naturally, independent of human activity
Alien (exotic, non-indigenous) species	A species that has been transported by human activity into an area outside its natural range
Introduced species	Alien species that has been transported by humans into an area outside its natural range, but has not yet established self-sustaining populations in the wild
Established (naturalised) species	Alien species that has been introduced and established self-sustaining populations in the wild
Feral species	Alien species that has been kept in captivity or domestication after introduction, but has escaped or been released to establish self-sustaining populations in the wild
Invasive species	Alien species that has been introduced, become established and is expanding its range, usually with deleterious consequences for native species
Introduction	The deliberate or accidental release into the wild of a non-native species
Translocation	The human-assisted movement of species within a specific region (i.e., movement of fish within rivers, lakes, ponds, etc.)
Dispersal	The natural dissemination (i.e., non-human assisted) of a species from its point of introduction

establish fishery resources.

Similar to trends observed in other countries, the number of aquatic invasive species in Iran has increased considerably over recent decades. These include harmful algal blooms (HAB) caused by *Cochlodinium polykrikoides* (Gymnodiniaceae) in the Persian Gulf in the south (Fatemi et al. 2012), *Mnemiopsis leidyi* blooms (MLB) in the Caspian Sea (Ivanov et al. 2000), *Azolla filiculoides* blooms (AFB) in Anzali wetland (Sadeghi et al. 2013) and invasion of the Oriental River Prawn, *Macrobrachium nipponense* in Anzali wetland (Grave & Ghane 2006) in the north, changing the species diversity and faunistic structure in Iranian water bodies. They have multiple synergistic impacts particularly on the freshwater ecosystem in this country through various ways.

Fishes are one of the aquatic groups which have been widely introduced and translocated. The movement of fishes between countries was reviewed by Welcomme (1981, 1988). Non-native species were primarily introduced into ecosystems in Iran through human activity, either deliberately or unintentionally and is well known, with the

introduction of *Gambusia* (Poeciliidae) in the 1920s and *Cyprinus carpio* (Cyprinidae) in the late 1930s when they were introduced to control malaria and to enhance fish production, respectively (Coad 1996; Esmaeili et al. 2007, 2010). Since then, about 30 fish species have been introduced or translocated and some of them are well established in different Iranian water bodies (Coad 1996; Esmaeili et al. 2007; 2010a, b, 2011, 2013; Khaefi 2014).

In this paper, we provide an update on diversity of alien and invasive fishes within Iranian freshwater ecosystems and discuss their possible environmental impacts. We additionally discuss the possible management strategies to protect the native (and endemic) ichthyodiversity in the 17 drainage basins of Iran recognized in Keivany & Ghorbani (2012), Keivany & Esmaeili (2013, 2014), and Keivany et al (2014).

Materials and Methods

This paper is mainly based on our field work data, since 2004 and also on Armantrout (1990), Holčík & Razavi (1991), Coad & Abdoli (1993), Coad (1995, 1996), Teimori (2006), Esmaeili et al. (2007,




Fig.1. Sympatric native and invasive species in Maharlo basin, except *Xiphophorus hellerii*. a, *Acanthobrama persidis*; b, *Cyprinus carpio*; c, *Capoeta damascina*; d, *Carasobarbus luteus*; e, *Ctenopharyngodon idella*; f, *Hypophthalmichthys molitrix*; g, *Pseudorasbora parva*; h, female *Aphanius farsicus*; i, male *Aphanius farsicus*; j, *Gambusia holbrooki*; k, *Xiphophorus hellerii*; l, *Chela abu*.

2010a,b, 2013, 2014), Hashemi & Ansary (2012), Hashemi et al. (2014), Keivany et al. (2014) and

Khaefi et al. (2014). Terminology given by Lymbery et al. (2014) and Gozlan et al. (2010) was followed

Table 2. Alien (exotic) and invasive freshwater fish species of Iran. E=exotic; *=invasive; NC= not confirmed in recent years.

Order	Family	Species	Status	Distribution
Anguilliformes	Anguillidae	<i>Anguilla anguilla</i>	E	Caspian Sea
		<i>Carassius auratus</i>	E*	Wide distribution
Cypriniformes	Cyprinidae	<i>Carassius carassius</i>	E, NC	Tigris and Caspian
		<i>Carassius gibelio</i>	E	Caspian Sea
		<i>Ctenopharyngodon idella</i>	E	Wide distribution
		<i>Cyprinus carpio</i>	E*	Wide distribution
		<i>Hemiculter leucisculus</i>	E	Caspian Sea/Urmia
		<i>Hypophthalmichthys molitrix</i>	E	Wide distribution
		<i>Hypophthalmichthys nobilis</i>	E	Wide distribution
		<i>Mylopharyngodon piceus</i>	E	Caspian Sea
		<i>Pimephales promelas</i>	E, NC	Namak Lake
		<i>Pseudorasbora parva</i>	E*	Wide distribution
Siluriformes	Heteropneustidae	<i>Heteropneustes fossilis</i>	E*	Tigris
Salmoniformes	Salmonidae	<i>Coregonus lavaretus</i>	E	Namak Lake
		<i>Oncorhynchus keta</i>	E	Caspian Sea
		<i>Oncorhynchus mykiss</i>	E	Wide distribution
		<i>Salmo trutta</i>	E	Caspian Sea
		<i>Salvelinus fontinalis</i>	E	Namak Lake
Mugiliformes	Mugilidae	<i>Chelon aurata</i>	E	Caspian Sea
		<i>Chelon saliens</i>	E	Caspian Sea
		<i>Mugil cephalus</i>	E	Caspian Sea
Beloniformes	Adrianichthyidae	<i>Oryzias latipes</i>	E, NC	Tedzhen
		<i>Gambusia holbrooki</i>	E *	Wide distribution
Cyprinodontiformes	Poeciliidae	<i>Gambusia affinis</i>	E, NC	?
		<i>Xiphophorus hellerii</i>	E*	Bushehr
		<i>Gasterosteus aculeatus</i>	E*	Caspian Sea
Gasterosteiformes	Gasterosteidae	<i>Rhinogobius similis</i>	E	Harirud
	Perciformes	Centrarchidae	<i>Lepomis macrochirus</i>	E, NC
<i>Micropterus salmoides</i>			E, NC	Tigris
Cichlidae		<i>Amatitlania nigrofasciata</i>	E	Hormuz
Pleuronectiformes	Pleuronectidae	<i>Tilapia zillii</i>	E	Tigris
		<i>Platichthys flesus</i>	E, NC	Caspian Sea

for alien species (Table 1).

Results

In the case of freshwater fishes, the number of reported alien and invasive fishes has reached about 32 species in 10 orders and 12 families (Table 2, Fig. 1) which is about 14.7% of the freshwater ichthyofauna of Iran. However, there are reports (e.g., Coad 1995) of some unconfirmed exotic and transplanted species such as *Carassius carassius* (Linnaeus, 1758), *Pimephales promelas* (Rafinesque, 1820), *Gambusia affinis* (Baird & Girard, 1853), *Oryzias latipes* (Temminck & Schlegel, 1846), *Lepomis macrochirus* (Rafinesque, 1819),

Micropterus salmoides (Lacèpede, 1802) and *Platichthys flesus* (Linnaeus, 1758). Hence, the confirmed exotic and invasive fishes comprise 25 species. Cyprinidae with 9 species (36% of confirmed non-native species) is ranked the first, and is followed by Salmonidae (5 species, 20%), Mugilidae (3 species, 12%), Cichlidae and Poeciliidae each with 2 confirmed species (8%), and 4 families each with only one species (4%). The Neotropical Convict cichlid, *Amatitlania nigrofasciata*, is a new alien fish species which is reported from Iran, the second record of this species from the Middle East and the only known extant wild population in the region (Esmaeili et al. 2013). The exotic cichlid, the

Table 3. Fishes of Shadegan Wetland. N= native; E= exotic; *= invasive.

Order	Family	Species	Status
Clupeiformes	Clupeidae	<i>Tenuulosa ilisha</i>	N
		<i>Sardinella sindensis</i>	N
	Engraulidae	<i>Thryssa hamiltonii</i>	N
Cypriniformes	Cyprinidae	<i>Acanthobrama marmid</i>	N
		<i>Alburnoides</i> sp.	N
		<i>Leuciscus vorax</i>	N
		<i>Carassius auratus</i>	E*
		<i>Carasobarbus luteus</i>	N
		<i>Chondrostoma regium</i>	N
		<i>Cyprinion kais</i>	N
		<i>Cyprinion macrostomum</i>	N
		<i>Ctenopharyngogon idella</i>	E
		<i>Cyprinus carpio</i>	E*
		<i>Hypophthalmichthys molitrix</i>	E
		<i>Hypophthalmichthys nobilis</i>	E
		<i>Luciobarbus barbulus</i>	N
		<i>Luciobarbus pectoralis</i>	N
		<i>Luciobarbus xanthopterus</i>	N
<i>Mesopotamichthys sharpeyi</i>	N		
<i>Tor grypus</i>	N		
Siluriformes	Heteropneustidae	<i>Heteropneustes fossilis</i>	E*
	Siluridae	<i>Silurus triostegus</i>	N
Mugiliformes	Mugilidae	<i>Chelon abu</i>	N
		<i>Chelon subviridis</i>	N
		<i>Ellohelon vaigiensis</i>	N
Cyprinodontiformes	Poeciliidae	<i>Gambusia holbrooki</i>	E*
Perciformes	Cichlidae	<i>Tilapia zillii</i>	E
	Sparidae	<i>Acanthopagrus arabicus</i>	N
Synbranchiformes	Mastacembelidae	<i>Mastacembelus mastacembelus</i>	N

redbelly tilapia, *Tilapia zillii* (Gervais, 1848), is another new record of an alien fish from natural freshwater bodies in Iran, which was most likely deliberately released to enhance fish production (Khaefi et al. 2014). Exotic and invasive species are distributed in all the 17 drainage basins of Iran and the Caspian Sea basin is ranked first in terms of invasive fish diversity (Table 2). Unfortunately some species have been introduced to fish hot spots of Iran including the Shadegan and Anzali wetlands in south and north of Iran, respectively changing the fish composition in these water bodies (Tables 3, 4). Shadegan wetland is one the largest international wetlands of Iran, situated in the southwest of the country and connecting the Jarahi River to the Persian Gulf. Due to its unique biodiversity, it has been considered as one the wonderful natural

landscapes of the world (Hashemi & Ansari 2012). Records of more than 29 fish species including 21 natives and 8 aliens show the high fish diversity of this wetland (Table 3). Introduction of *Tilapia zillii* to this international wetland may have sever affects on native fishes.


Anzali wetland with about 50 native fish species is one of the fish hot spots of Iran. However, during the past few decades about 16 alien fish species (Table 4) have been introduced to this water body changing its fish composition and affecting native fishes. Moreover, additional invaders such as blooms of *Azolla filiculoides* (Sadeghi et al. 2013) and invasion of the Oriental River Prawn, *Macrobrachium nipponense* (Grave & Ghane 2006) in Anzali wetland could have multiple synergistic impacts on this unique freshwater ecosystem.

Table 4. Fishes of Anzali Wetland. N= native; E= exotic; *= invasive.

Order	Family	Species	Status
Petromyzontiformes	Petromyzontidae	<i>Caspiomyzon wagneri</i>	N
Acipenseriformes	Acipenseridae	<i>Acipenser persicus</i>	N
		<i>Acipenser stellatus</i>	N
Anguilliformes	Anguillidae	<i>Anguilla anguilla</i>	E
Clupeiformes	Clupeidae	<i>Alosa brashnikovi</i>	N
		<i>Alosa caspia</i>	N
		<i>Alosa knipowitschia</i>	N
		<i>Alosa persica</i>	N
		<i>Alosa kessleri</i>	N
		<i>Clupeonella caspia</i>	N
		Cypriniformes	Cyprinidae
<i>Alburnoides eichwaldii</i>	N		
<i>Alburnus chalcoides</i>	N		
<i>Alburnus filippii</i>	N		
<i>Alburnus hohenackeri</i>	N		
<i>Leuciscus aspius</i>	N		
<i>Barbus cyri</i>	N		
<i>Blicca bjoerkna</i>	N		
<i>Capoeta capoeta</i>	N		
<i>Carassius auratus</i>	E*		
<i>Carassius gibelio</i>	E*		
<i>Ctenopharyngodon idella</i>	E		
<i>Cyprinus carpio</i>	E*		
<i>Hemiculter leucisculus</i>	E		
<i>Hypophthalmichthys molitrix</i>	E		
<i>Hypophthalmichthys nobilis</i>	E		
<i>Leucaspis delineatus</i>	N		
<i>Luciobarbus brachycephalus</i>	N		
<i>Luciobarbus capito</i>	N		
<i>Mylopharyngodon piceus</i>	E		
<i>Pelecus cultratus</i>	N		
<i>Pseudorasbora parva</i>	E		
<i>Rhodeus amarus</i>	N		
<i>Rutilus kutum</i>	N		
<i>Rutilus caspicus</i>	N		
<i>Scardinius erythrophthalmus</i>	N		
<i>Squalius cephalus</i>	N		
<i>Tinca tinca</i>	N		
<i>Vimba persa</i>	N		
<i>Cobitis faridpaki</i>	N		
<i>Cobitis keyvani</i>	N		
Cobitidae	<i>Sabanejewia aurata</i>	N	
	<i>Sabanejewia caspia</i>	N	
	<i>Oxynoemacheilus bergianus</i>	N	
Nemacheilidae	Nemacheilidae	<i>Oxynoemacheilus bergianus</i>	N
		<i>Silurus glanis</i>	N
Siluriformes	Siluridae	<i>Silurus glanis</i>	N
Salmoniformes	Salmonidae	<i>Oncorhynchus mykiss</i>	E
		<i>Salmo caspius</i>	N
		<i>Salmo fario</i>	N
Esociformes	Esocidae	<i>Esox lucius</i>	N
Mugiliformes	Mugilidae	<i>Chelon aurata</i>	E
		<i>Chelon saliens</i>	E

Table 4. Continued

Order	Family	Species	Status
Atheriniformes	Atherinidae	<i>Atherina caspia</i>	N
Cyprinodontiformes	Poeciliidae	<i>Gambusia holbrooki</i>	E*
Gasterosteiformes	Gasterosteidae	<i>Gasterosteus aculeatus</i>	E*
		<i>Pungitius platygaster</i>	N
Syngnathiformes	Syngnathidae	<i>Syngnathus caspius</i>	N
Perciformes	Percidae	<i>Perca fluviatilis</i>	N
		<i>Sander lucioperca</i>	N
		<i>Benthophilus kessleri</i>	N
	Gobiidae	<i>Knipowitschia caucasica</i>	N
		<i>Neogobius melanostomus</i>	N
		<i>Ponticola cyrius</i>	N
		<i>Proterorhinus nasalis</i>	N
		<i>Rhinogobius</i> sp.	N

**Fig.2.** Schematic diagram of processes involved in species invasions (Modified from Lymbery et al. 2014).

The Eastern Mosquitofish, *Gambusia holbrooki* is the most widespread invasive species while *A. nigrofasciata* (in Hormuz basin) and *Xiphophorus hellerii* (Persian Gulf basin) have restricted distributions. Some of the invasive species (e.g., *Cyprinus carpio*, *Carassius auratus*, *Pseudorasbora parva*, *Xiphophorus hellerii* and *Gambusia*

holbrooki) have established breeding populations, acting as invaders, while some others are regularly stocked by the Fisheries Department of Iran (e.g., *Hypophthalmichthys molitrix*, *H. nobilis*, *Ctenopharyngodon idella*) and a few others are occasionally recorded from natural waters.

The processes involved in species invasions are




Fig.3. Schematic diagram of ecological impacts of nonnative freshwater fishes. (Modified from Cucherousset & Olden 2011).

almost similar in every biological invasion (Fig. 2). An alien species must surmount geographic barriers to be introduced into a new area, then barriers to survival and reproduction, to become established within the expanded range, and finally barriers to dispersal to become invasive (Blackburn et al. 2011; Lymbery et al. 2014). Due to the nature of the neotropical convict cichlid, *Amatitlania nigrofasciata* and the redbelly tilapia, *Tilapia zillii*, they also may overcome this common framework and act as invasive species in near future.

Discussion

Similar to trends observed in other world regions, the number of non-native fish species introduced and established in Iran has increased considerably over recent decades attaining to 25 confirmed species comprising about 11.5% of the total confirmed freshwater fishes in this country. Aquaculture, sport fishing, control of malaria, ornamental purposes, research activities, demonstration in national fairs

and accidental introduction are the main reasons for these introductions (Coad & Abdoli 1993; Esmaeili et al. 2010a, b). Some of them (e.g., *Cyprinus carpio*, *Carassius auratus*, *Pseudorasbora parva*, *Xiphophorus hellerii* and *Gambusia holbrooki*) have been established in natural water bodies acting as invasion species. Fish behavior, minimal resource competition, suitable water temperature, changes in water flow, suitable habitat for spawning, number of release events, species-level factors (such as diet, offspring per year, growth rate, body size, lifespan) and abundant food supply have been recorded to be the main factors promoting establishment of introduced fish species. Whether through accidental introductions or intentional releases, many species attain distributions beyond their natural ranges and form novel interactions with native counterparts (Paterson et al. 2013). Introductions always carry risks for the native biota. The introduction of a non-native species in an ecosystem is always likely to present an ecological risk if the species is able to integrate itself successfully into the ecosystem (Gozlan & Newton 2009), resulting in possible detrimental interactions with native species or even on ecosystem functioning (Gozlan et al. 2010). The ecological consequences of freshwater fish invasions have been reviewed and discussed by Cucherousset & Olden (2011) which are given in Fig. 3. The effects of invaders could be at genetic, individual, population, community and ecosystem levels.

Much of what we know about invaders is based on information on introduced populations in other countries. Results of these past introductions provide perspective on possible outcomes for Iran. The following parameters suggest being the most important impacts of the invasive species on Iranian freshwater fishes (native and endemic).

Competition: Mosquito fish (*Gambusia holbrooki*) prey on the eggs of others fishes; they are aggressive, even attacking fish larger than themselves, thus discouraging feeding and reproduction, and compete directly for food with various cyprinid species, so the mosquito fish has been called "the fish destroyer" by

Myers (1965) and is said to replace native species aggressively (Armantrout 1990). In Iran, the native *Aphanius* species occurs alongside *Gambusia* wherever the latter has been introduced. *Pseudorasbora parva*, competes strongly with other fishes for food and is a predator on their eggs and young.

Effects of redbelly tilapia on native organisms have been well documented. Molnar et al. (2008) considered it as a potential competitor with native fish for food and spawning areas, and it is potentially detrimental to California rice crops. This voracious herbivore constitutes a serious threat to native aquatic plants and the organisms that depend on them. Courtenay in Hogg (1976) called it the most destructive fish to submerged vegetation known next to the grass carp, *Ctenopharyngodon idella*. Due to its feeding habits, the invasive redbelly tilapia eliminated all aquatic macrophytes from the Hyco Reservoir, North Carolina, USA, within a 2-year period, which caused declines in populations of several native fishes; however, because of its ability to switch to alternate food sources, the tilapia populations continued expanding in the absence of macrophytes (Molnar et al. 2008). *Amatitlania nigrofasciata* may cause harm to native fishes as this is an aggressive and territorial species which is well known for its ability to colonize rapidly and it also often carries alien parasites (Wisenden 1994; Bassleer 1997; Martinez et al. 2002).

Habitat change: Some introduced species change the fish habitat. *Gambusia holbrooki* has been shown to have major effects on habitats under experimental conditions (Hurlbert et al. 1972). Grass carp, *Ctenopharyngodon idella* removes the habitat vegetation and decreases the area of refuge for larvae and smaller fish and production of invertebrates on which many native species feed. Carp and goldfish, feed on the bottom and in so doing muddy water.

Genetic changes: Many native species exist in several

water bodies in a desert environment and the systematic status of these isolated populations has not been examined (Coad 1995). Movement of taxa between these water bodies may well eliminate a diversity as yet unrevealed. Populations of the bighead (*Hypophthalmichthys nobilis*) in Anzali Wetland of Iran are hybrids with silver carp (*Hypophthalmichthys molitrix*) (Holcik 1989). Hybrids of *Cyprinus carpio* and *Carassius auratus* have been observed in the Caspian Sea basin by K. Abbasi.

Introduction of parasites: Introduced fish species could play a role in the spread of disease or parasites to other animals. As many parasites have been recorded from invasive and native fish species of Iran (see Coad 2014), it seems that the wide distribution range of invasive species may have a significant role in the spread of these parasites. Aquatic invasive species (AIS) have multiple synergistic impacts on the ecosystem through various ways of hybridization, gene pollution, competition, transmission of pathogens and parasites, community level changes, facilitating the introduction of other taxa and elimination and subsequent extinction of endemic species. It can be considerably critical when the invasive species are sympatric to a native (and endemic) species, as it can be seen in the case of several endemic fishes in Iran. A particular example is the cyprinodontid fishes (see above).

Management: The intentional introduction of exotic species is an extraordinarily complex issue that requires taking into account a broad number of variables. Understanding the risks that alien species, especially aquatic invasive species (AIS) pose, requires comparative studies on the ecology and life history strategies of invasive and closely related indigenous species with similar ecological life traits, correct identification of AIS, monitoring the rate of invasion, the causal relation between population dynamics and species invasion, regular checking of

fishes for diseases and parasites, obtaining a more balanced perspective on the ecological impacts of fish invasions across levels of biological organization, improving our understanding of how other facets of environmental change (e.g., habitat degradation, land-use change, climate warming) interact with invasive species, elucidation of the adaptive and evolutionary responses of native species to the presence of invasive species and vice versa (see Cucherousset & Olden 2011), and public awareness, all of which could be effective management strategies to minimize the bio-invasion impacts. Priority may be given to those invasive species that have already been established in the wild and on preventing new infestations.

Finally, as suggested by Gozlan et al. (2010) maps of conservation hot spots should be drawn based on a combination of predictors of future introductions (e.g., human population density, aquaculture activity) and risk level of incurring losses to local– global biodiversity (e.g., number of critically endangered species). This would accelerate prioritization of the areas at national and global levels where risks of future introductions should be minimized, such as those where endemic species are at risk and the intensity of introductions of non-native species has been limited.

Acknowledgements

We would like to thank H.R. Mehraban and A. Gholamifard for helping with the drawings and photography. This study was financially supported by Shiraz University.

References

Armantrout, N.B. 1990. Fish distribution as influenced by aquatic habitat alteration and species introductions, In: Daniel, J.C. & Serrao, J.S. (Eds). Conservation in developing countries: problems and prospects. Proceedings of the Centenary Seminar of the Bombay Natural History Society. Bombay, Delhi, Calcutta, Madras. pp. 199-206.

Bassleer, G. 1997. Color Guide of Tropical Fish Diseases:

on Freshwater Fish. Bassleer Biofish, Westmeerbeek, Belgium.

Blackburn, T.M.; Pyšek, P.; Bacher, S.; Carlton, J.T.; Duncan, R.P.; Jarošík, V.; Wilson, J.R.U. & Richardson, D.M. 2011. A proposed unified framework for biological invasions. *Trends in Ecology and Evolution* 26: 333–339.

Coad, B.W. 1995. Freshwater Fishes of Iran. *Acta Scientiarum Naturalium Academiae Scientiarum Bohemicae*, Brno, 29(1): 1-64.

Coad, B.W. 1996. Exotic and transplanted fishes in Southwest Asia. *Publicaciones Especiales Instituto Español de Oceanografía* 21: 81-106.

Coad, B.W. 2014. Freshwater Fishes of Iran. www.briancoad.com.

Coad, B.W. & Abdoli, A. 1993. Exotic fish species in the freshwaters of Iran. *Zoology in the Middle East* 9: 65-80.

Cucherousset, J. & Olden, J.D. 2011. Ecological Impacts of Non-native Freshwater Fishes. *Fisheries* 36(5): 215-230.

Esmaeili, H.R.; Teimori, A. & Gholamhosseini, G. 2007. Freshwater Ichthyodiversity and its conservation in Iran. Xii European Congress on Ichthyology. Cavata (Dubrovnik), Croatia, pp. 200-201.

Esmaeili, H.R.; Coad, B.W.; Gholamifard, A.; Nazari, N. & Teimori, A. 2010a. Annotated checklist of the freshwater fishes of Iran. *Zoosystematica Rossica* 19: 361–386.

Esmaeili, H.R.; Gholamifard, A.; Teimori, A.; Bagh-Bani, S. & Coad, B.W. 2010b. *Xiphophorus hellerii* Heckel, 1848 (Cyprinodontiformes, Poeciliidae), a newly introduced fish recorded from natural freshwaters of Iran. *Journal of Applied Ichthyology* 26: 937-938.

Esmaeili, H.R.; Nazari, N.; Gholamifard, A.; Gholamhosseini, G.; Teimori, A. & Coad, B.W. 2011. Range extension and translocation for *Rhodeus amarus* (Bloch, 1782) Actinopterygii: Cyprinidae) in northwest Iran. *Turkish Journal of Zoology* 35(6): 883-886.

Esmaeili, H.R.; Gholamifard, A.; Sayyadzadeh, G.; Parsi, B.; Mirghiyasi, S. & Ghasemian, S. 2013. New record of the convict cichlid, *Amatitlania nigrofasciata* (Günther, 1867), from the Middle East (Actinopterygii: Cichlidae). *Aqua* 19: 225-229.

- Esmaeili, H.R.; Masoudi, M. & Mehraban, H.R. 2014. Assignment of *Acanthopagrus populations* in the Persian Gulf drainage system of Iran to *Acanthopagrus arabicus* Iwatsuki, 2013 (Perciformes: Sparidae). Iranian Journal of Ichthyology 1(1): 23-28.
- Fatemi, S.M.R.; Nabavi, S.M.B.; Vosoghi, G.; Fallahi, M. & Mohammadi, M. 2012. The relation between environmental parameters of Hormuzgan coastline in Persian Gulf and occurrence of the first harmful algal bloom of *Cochlodinium polykrikoides* (Gymnodiniaceae). Iranian Journal of Fisheries Sciences 11(3): 475-489.
- Gozlan, R.E. & Newton, A.C. 2009. Biological invasions: benefits versus risks. Science 324: 1015-1016.
- Gozlan, R.E.; Britton, J.R.; Cowx, I. & Copp, G.H. 2010. Current knowledge on non-native freshwater fish introductions. Journal of Fish Biology 76: 751-786.
- Grave, S.D. & Ghane, A. 2006. The establishment of the Oriental River Prawn, *Macrobrachium nipponense* (de Haan, 1849) in Anzali Lagoon, Iran. Aquatic Invasions 1 (4): 204-208.
- Hashemi, S.A.R. & Ansary, H. 2012. Biomass and production of fish species in the Shadegan wetland, Iran. Research Journal of Recent Science 1: 66-68.
- Hashemi, S.A.R.; Ghorbani, R.; Kymaram, F.; Hossini, S.A.; Eskandari, G. & Hedayati, A. 2014. Estimation of fish composition and catchability coefficient of gillnet in the Shadegan Wetland. Iranian Journal of Ichthyology 1(1): 51-60.
- Hogg, R.G. 1976. Established exotic cichlid fishes in Dade County, Florida. Florida Scientist 39(2): 97-103.
- Holčík, J. 1989. The Freshwater Fishes of Europe. Volume I, Part II. General Introduction to Fishes. Acipenseriformes. AULA-Verlag, Wiesbaden.
- Holčík, J. & Razavi, B. A. 1991. New species of freshwater fish from the Iranian coast of the Caspian Sea. Report prepared for the Anzali Lagoon productivity and fish stocks investigation project. Food and Agriculture Organization, Rome, FI/UNDP/IRA/88/001 Field Document 1. En, 14 pp.
- Hurlbert, S., Zeidler, J. & Fairbanks, D. 1972. Ecosystem alteration by mosquito fish (*Gambusia affinis*) predation. Science 175: 639-641.
- Ivanov, V.P.; Kamakin, A.M.; Ushivtzev, V.B.; Shiganova, T.; Zhukova, O.; Aladin, N.; Wilson, S.I.; Harbison, G.R. & Dumont, H.J. 2000. Invasion of the Caspian Sea by the comb jellyfish *Mnemiopsis leidyi* (Ctenophora). Biological Invasions 2: 255-258.
- Keivany, Y. & Ghorbani, M. 2012. Distribution of *Aphanius dispar dispar* (Rüppell, 1829) populations in Iran, with a new record from western Iran (Actinopterygii: Cyprinodontidae). Turkish Journal of Zoology 36(6): 824-827.
- Keivany, Y. & Esmaeili, H.R. 2013. Threatened fishes of the world: *Aphanius farsicus* Teimori, Esmaeili & Reichenbacher, 2011 (Cyprinodontidae). Croatian Journal of Fisheries 71(4): 192-194.
- Keivany, Y. & Esmaeili, H.R. 2014. Threatened fishes of the world: *Aphanius pluristriatus* (Jenkins, 1910) (Cyprinodontidae). Aqua 20(2): 67-72.
- Keivany, Y.; Nasri, M.; Abbasi, K. & Abdoli, A. 2014. Atlas of inland water fishes of Iran. Iran Department of Environment Press. In press.
- Khaefi, R.; Esmaeili, H.R.; Zareian, H. & Babaei, S. 2014. The first record of the redbelly tilapia, *Tilapia zillii* (Gervais, 1848), in freshwaters of Iran. Turkish Journal of Zoology 38: 96-98.
- Kolar, C.S. & Lodge, D.M. 2001. Progress in invasion biology: predicting invaders. Trends in Ecology and Evolution. 16: 199-204.
- Lymbery, A.J.; Morine, M.; Kanani, H.J.; Beatty, S.J. & Morgan, D.J. 2014. Co-invaders: The effects of alien parasites on native hosts. International Journal for Parasitology: Parasites and Wildlife. Available online 24 April 2014. <http://dx.doi.org/10.1016/j.ijppaw.2014.04.002>.
- Martinez, V.M.V.; Macedo, M.L.A.; Scholz, T.; Solis, D.G. & Franco, E. F. M. 2002. Atlas de los helmintos parásitos de cíclidos de México. Instituto Politécnico Nacional, Dirección de Publicaciones Tresguerras.
- Molnar, J.L.; Gamboa, R.L.; Revenga, C. & Spalding, M.D. 2008. Assessing the global threat of invasive species to marine biodiversity. Frontiers in Ecology and the Environment 6: 485-492.
- Myers, G.S. 1965. Gambusia, the fish destroyer. Tropical Fish Hobbyist 13(5): 31-32, 53-54.
- Paterson, R.A.; Lal, A.; Dale, M.; Townsend, C.R.;

- Poulin, R. & Tompkins, D.M. 2013. Relative competence of native and exotic fish hosts for two generalist native trematodes. *International Journal for Parasitology: Parasites and Wildlife* 2: 136–143.
- Sadeghi, R.; Zarkami, R.; Sabetraftar, K. & Damme, P. V. 2013. A review of some ecological factors affecting the growth of *Azolla* spp. *Caspian Journal of Environmental Sciences* 1 (1): 65-76.
- Sakai, A.K.; Allendorf, F.W.; Holt, J.S.; Lodge, D.M.; Molofsky, J.; With, K.A.; Baughman, S.; Cabin, R.J.; Cohen, J.E.; Ellstrand, N.C.; McCauley, D.E.; O’Neil, P.; Parker, I.M.; Thompson, J.N. & Weller, S.G. 2001. The population biology of invasive species. *Annual Review of Ecology and Systematics* 32: 305–332.
- Teimori, A. 2006. A preliminary study of freshwater fish diversity in southern Iran, Fars province. Master Thesis. Department of Biology, Shiraz University, Shiraz, Iran.
- Vitousek, P.M.; Mooney, H.A.; Lubchenko, J. & Melillo, J.M. 1997. Human domination of Earth’s ecosystems. *Science* 277: 494–499.
- Welcomme, R.L. 1981. Register of International Transfers of Inland Fish Species. Food and Agriculture Organization, Rome, Fisheries Technical Paper 213.
- Welcomme, R.L. 1988. International introductions of inland aquatic species. Food and Agriculture Organization, Rome, Fisheries Technical Paper 294.
- Wisenden, B.D. 1994. Factors affecting mate desertion by males in free-ranging convict cichlids (*Cichlasoma nigrofasciatum*). *Behavioral Ecology* 5(4): 439-447.